

2017 CMS Quality Reporting - ABSTRACTION PROCESS OVERVIEW

REMEMBER! Abstract only those cases belonging to your ACO and use only MiShare (<https://mishare.med.umich.edu/>) when communicating PHI!

STEP 1: Can you locate the patient's medical record?

YES

NO

STEP 2: Does 1 or more of the following 4 conditions apply to the patient?

Select "No - Medical Record Not Found" and proceed to next patient in queue

- **Deceased** - patient died **at any time** during the measurement period
- **In hospice** - patient was in hospice care **at any time** during the measurement period, including non-hospice patients receiving palliative goals or comfort care
- **Moved out-of-the-country** - patient moved out of the country **any time** during the measurement period
- **Enrolled in HMO** - patient enrolled in HMO **at any time** during the measurement period, i.e., Medicare Advantage, non-Medicare HMOs, etc.

NO

STEP 3: Is there another CMS-approved reason for disqualification?

NOTE: To request a CMS Approved Reason, you need to open a Quality Payment Program Service Inquiry, and provide the following information:

- patient rank
- title of the measure in question
- reason for request

YES

YES

NO

- (1) Select "Not Qualified for Sample"
- (2) Select the drop-down **reason** for not qualifying that applies to the case in question
- (3) Enter the **date** when the patient became ineligible
 - If the date that the patient became ineligible is unknown, enter the last day of the measurement period, i.e., 12/31/17

- Enter the Help Desk ticket number for the approved reason in the "Help Desk Ticket #" field located within the Web Interface
- The same Help Desk ticket number can be used for multiple patients with identical reasons
- Proceed to next patient or measure in the queue

STEP 4: Is there a measure-specific reason for disqualification?

➤ See **MEASURE-SPECIFIC EXCLUSIONS AND REQUIRED DOCUMENTATION** (pgs. 2-4) for additional, measure-specific exclusions and **required documentation** for same

CONGRATULATIONS! You're ready to abstract!

NO

2017 CMS Quality Reporting - MEASURE-SPECIFIC EXCLUSIONS AND REQUIRED DOCUMENTATION

Measure	Measure Title	Measure-Specific Exclusions
CARE-1	Medication Reconciliation Post-Discharge	<p>No current medications listed</p> <p>No discharge medications listed</p> <p>No follow-up visit in chart</p> <p>Follow-up > 32 days (<i>Rule states 30 days, +/- 2 days from prefilled discharge date</i>)</p> <p>Author is not a prescribing practitioner, RN, or clinical pharmacist</p>
CARE-2	Falls - Screening for Future Falls	<p>Patient non-ambulatory at most recent encounter on 9/1/17</p>
DM-2	Diabetes - Hemoglobin A1c (HbA1c) Poor Control (>9%)	<p>Patient has no diagnosis of Type 1 or 2 diabetes on chart</p> <p>Patient has no <i>active</i> diagnosis of Type 1 or 2 diabetes on chart</p> <p>Elevated blood sugar is due to another medical condition, e.g., steroids for asthma</p> <p>Hemoglobin A1c (HbA1c) is ≤ 9%</p> <p>No Hemoglobin A1c (HbA1c) on chart</p>
DM-7	Diabetes - Eye Exam	<p>Patient has no diagnosis of Type 1 or 2 diabetes on chart</p> <p>Patient has no <i>active</i> diagnosis of Type 1 or 2 diabetes on chart during MP</p> <p>Elevated blood sugar is due to another medical condition, e.g., steroids for asthma</p> <p>Retinal or dilated eye exam was not performed as part of exam</p> <p>No eye exam on chart</p> <p>Exam performed outside of measure-specified time period, i.e., 1/1/2016 - 12/31/2017</p>
HTN-2	Controlling High Blood Pressure	<p>No active diagnosis of essential hypertension between 01/01 and 06/30/2017</p> <p>Systolic blood pressure > 140</p> <p>Diastolic blood pressure > 90</p>
IVD-2	Ischemic Vascular Disease (IVD) - Use of Aspirin or Another Antiplatelet	<p>No acute myocardial infarction between 01/01 - 12/31/2016</p> <p>No coronary artery bypass graft between 01/01 - 12/31/2016</p> <p>No percutaneous coronary intervention between 01/01 - 12/31/2016</p> <p>No active diagnosis of ischemic vascular disease during the measurement period</p> <p>No documented use of aspirin or other antiplatelet medication during the measurement period</p> <p>Patient is taking one of the Denominator Exclusion Anticoagulant Medications</p>
MH-1	Depression Remission at Twelve Months	<p>Patient has an active diagnosis of bipolar disorder</p> <p>Patient has an active diagnosis of personality disorder</p> <p>No <u>index</u> PHQ-9 on chart between 12/01/2015 - 11/30/2016</p> <p>Index PHQ-9 score is ≤ 9</p> <p>No <u>repeat</u> PHQ-9 on chart at 12 months after the <u>index</u> PHQ-9 (+/- 30 days)</p> <p>Repeat PHQ-9 score ≥ 5</p>

2017 CMS Quality Reporting - MEASURE-SPECIFIC EXCLUSIONS AND REQUIRED DOCUMENTATION

Measure	Measure Title	Measure-Specific Exclusions
PREV-5	Breast Cancer Screening	<p>Patient had a bilateral mastectomy</p> <p>Patient had two (2) unilateral mastectomies</p> <p>No mammogram on chart between 10/01/2015 - 12/31/2017</p> <p>Most recent mammogram was before 10/01/2015</p> <p>Patient received an MRI or Ultrasound instead of a mammogram</p> <p>Patient-reported results are listed by the provider, however either the test type, date, or result is missing</p>
PREV-6	Colorectal Cancer Screening	<p>Patient has a history of colorectal cancer</p> <p>Patient has a history of total colectomy</p> <p>Patient received only digital rectal exam</p> <p>Patient-reported results are listed by the provider, however either the test type, date, or result is missing</p>
PREV-7	Influenza Immunization	<p>No chart documentation reflecting that the patient received an influenza immunization during a clinic visit between 10/01/2016 - 03/31/2017</p> <p>No chart documentation reflecting that the patient reported having previously received an influenza immunization between 08/01/2016 - 03/31/2017</p>
PREV-8	Pneumococcal Vaccination Status for Older Adults	<p>No chart documentation that the patient ever received either the PCV13 or PPSV23 pneumonia vaccination, either alone or in combination</p> <p>Patient-reported results are listed by the provider, however either the test type, date, or result is missing</p>
PREV-9	Body Mass Index (BMI) Screening and Follow-Up Plan	<p>Patient is pregnant</p> <p>Patient refused height and/or weight measurement</p> <p>Patient refused follow-up</p> <p>Patient is ≥ 65 years and weight loss or gain would complicate other health conditions, such as: illness or physical disability; mental illness, dementia, confusion; or vitamin/mineral deficiency</p> <p>Patient is emergent or urgent and a treatment delay would jeopardize patient health</p> <p>No BMI charted during current encounter or during the six (6) months prior to current encounter</p> <p>Patient's Body Mass Index is on the patient chart and is within normal range</p> <p>Patient's Body Mass Index is on the chart and abnormal but there is no follow-up plan</p>
PREV-10	Tobacco Use - Screening and Cessation Intervention	<p>Patient was not screened between 01/01/2016 and 12/31/2017</p> <p>Patient is a non-smoker</p> <p>Patient had a positive screen however cessation counseling was not provided to the patient</p>

2017 CMS Quality Reporting - MEASURE-SPECIFIC EXCLUSIONS AND REQUIRED DOCUMENTATION

Measure	Measure Title	Measure-Specific Exclusions
PREV-12	<i>Screening for Depression and Follow-Up Plan</i>	Patient has an active diagnosis of bipolar disorder Patient has an active diagnosis of personality disorder No depression screen on chart Patient refuses assessment Patient is emergent or urgent and a treatment delay would jeopardize patient health Patient's functional capacity or motivation to improve may impact the accuracy of results Patient was not screened with a standardized tool Patient has positive screen but no follow-up plan is documented Follow-up plan is present but missing the minimum requirement Follow-up plan does not refer to the positive screen
PREV-13	<i>Statin Therapy for the Prevention and Treatment of Cardiovascular Disease</i>	Patient not "high-risk" i.e., patient does not meet at least one (1) of the three (3) reporting criteria for this measure Patient pregnant or breastfeeding or diagnosed with rhabdomyolysis Patient is allergic or intolerant of statin medication Patient has active liver/hepatic disease or insufficiency Patient has end-stage renal disease (ESRD) Patient is diabetic with most recent fasting or direct LDL-C < 70 mg/dl and is not taking a statin

2017 CMS Quality Reporting Quick Reference Guide

Measure	Description	Exclusions	Definitions	Data Guidance	
CARE-1 Medication Reconciliation Post-Discharge	<p>Patients ≥ 18 yrs who were:</p> <ul style="list-style-type: none"> -- discharged from an inpatient facility <li style="text-align: center;">AND -- seen within 30 days post-discharge (+ or - 2 days either before or after the pre-filled discharge date) <li style="text-align: center;">BY -- a professional providing ongoing care <li style="text-align: center;">AND -- discharge medication list was reconciled <li style="text-align: center;">WITH -- the current medication list in the MR. 	<ul style="list-style-type: none"> • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>Inpatient facility</i> = acute care hospital, inpatient psychiatric facility, skilled nursing or inpatient rehabilitation • <i>HMO</i> = Health Maintenance Organization • <i>MP</i> = Measurement Period = 01/01/2017 - 12/31/2017 • <i>MR</i> = Medical Record • <i>OV</i> = Office visit • <i>Professional providing ongoing care</i> = physician, prescribing practitioner, registered nurse, or clinical pharmacist • <i>WI</i> = Web Interface 	<ol style="list-style-type: none"> (1) Inpatient discharges are pre-populated by CMS using claims data (2) Denominator is based on individual discharges followed by an OV, not individual patients. As a result, patients may appear multiple times. (3) Documentation of the current medications with a notation that references the discharge medications, such as: <ul style="list-style-type: none"> -- <i>no changes in meds since discharge</i> -- <i>same meds at discharge</i> -- <i>discontinue all discharge meds</i> (4) Documentation of the patient's current medications with a notation that the discharge medications were reviewed. (5) Documentation that the provider "reconciled the current and discharge meds." (6) Documentation of a current medication list, a discharge medication list <u>and</u> notation that the appropriate practitioner type reviewed both lists on the same date of service (7) Notation that no medications were prescribed or ordered upon discharge. (8) Medication reconciliation post-discharge <u>can</u> be performed during telehealth encounter 	
CARE-2 Falls - Screening for Future Fall Risk	<p>Patients ≥ 65 yrs who were:</p> <ul style="list-style-type: none"> -- screened for future fall risk during the MP 	<ul style="list-style-type: none"> • Patient non-ambulatory <i>at most recent encounter during MP</i> • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>Fall</i> = A sudden, unintentional change in position causing an individual to land at a lower level, on an object, the floor, or the ground, other than as a consequence of sudden onset of paralysis, epileptic seizure, or overwhelming external force. • <i>HMO</i> = Health Maintenance Organization • <i>MP</i> = Measurement Period = 01/01/2017 - 12/31/2017 • <i>MR</i> = Medical Record • <i>Screening for Future Fall Risk</i> = assessment of whether an individual has experienced a fall or problems with gait or balance, e.g., Morse Fall Scale, timed Get-Up-And-Go test, etc. • <i>WI</i> = Web Interface 	<ol style="list-style-type: none"> (1) A clinician with appropriate skills and experience may perform the screening (2) <i>Documentation of no falls is sufficient</i> (3) MR must include documentation of screening performed (4) Any history of falls screening during the MP is acceptable (5) A gait or balance assessment is acceptable (6) A specific screening tool is not required (7) Fall risk screening is embedded within the Medicare Annual Wellness Visit documentaion (8) Fall risk screening <u>can</u> be performed during telehealth encounter 	
DM Composite - All or Nothing	DM-2 Diabetes - Hemoglobin A1c (HbA1c) Poor Control (>9%)	<p>Patients 18 - 75 yrs during MP with:</p> <ul style="list-style-type: none"> -- active diagnosis or documented history of Type 1 or Type 2 DM during MP or year prior to MP <li style="text-align: center;">AND -- <i>most recent</i> HbA1c > 9% during the MP 	<ul style="list-style-type: none"> • Diagnosis of secondary DM due to other condition • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>Active Diagnosis</i> = a diagnosis that is: <ul style="list-style-type: none"> -- on the patient's problem list -- a diagnosis code on the encounter, or -- documented in a progress note indicating that the patient is being treated or managed for the disease or condition during the MP • <i>DM</i> = Diabetes Mellitus • <i>HMO</i> = Health Maintenance Organization • <i>MP</i> = Measurement Period = 01/01/2017 - 12/31/2017 • <i>MR</i> = Medical Record • <i>WI</i> = Web Interface 	<ol style="list-style-type: none"> (1) Synonyms for HbA1c testing may include Glycohemoglobin A1c, HbA1c, Hemoglobin A1c, HgbA1c, A1c (2) Data source priority ranking: <ul style="list-style-type: none"> -- Lab report draw date (also known as collection date) -- Lab report date -- Flow sheet documentation -- Practitioner notes -- Other documentation -- Patient reported result (source of last resort) (3) Both patient reported and chart listed results must include date of test and numeric result (range unacceptable) (4) Documentation of <i>most recent</i> HbA1c result <u>can</u> be completed during a telehealth encounter
	DM-7 Diabetes - Eye Exam	<p>Patients 18 - 75 yrs with:</p> <ul style="list-style-type: none"> -- active diagnosis or documented history of Type 1 or Type 2 DM during MP or year prior to MP and -- a retinal or dilated eye exam performed or read by -- an ophthalmologist or optometrist during the MP <li style="text-align: center;">OR -- a neg. retinal or dilated eye exam (e.g. no evidence of retinopathy) within -- the 12 months <u>prior</u> to the MP 	<ul style="list-style-type: none"> • Diagnosis of secondary DM due to other condition • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>Active Diagnosis</i> = a diagnosis that is: <ul style="list-style-type: none"> -- on the patient's problem list -- a diagnosis code on the encounter, or -- documented in a progress note indicating that the patient is being treated or managed for the disease or condition during the MP • <i>DM</i> = Diabetes Mellitus • <i>HMO</i> = Health Maintenance Organization • <i>MP</i> = Measurement Period = 01/01/2017 - 12/31/2017 • <i>MR</i> = Medical Record • <i>WI</i> = Web Interface 	<ol style="list-style-type: none"> (1) Patient Reported Requirement: Date (year) and result/finding (2) If an endocrinologist or PCP performs the appropriate imaging in their office and the results are reviewed by an eye care professional (<i>optometrist or ophthalmologist</i>) during the MP or the year prior to the MP (if negative for retinopathy) then it is eligible for use in reporting (3) If the eye exam is not performed or reviewed by an ophthalmologist or optometrist, results may be read by a qualified reading center that operates under the direction of a medical director who is a retinal specialist (4) Documentation of diabetic retinal disease screening <u>can</u> be completed during a telehealth encounter

2017 CMS Quality Reporting Quick Reference Guide

Measure	Description	Exclusions	Definitions	Data Guidance
HTN-2 Controlling High Blood Pressure	Patients 18 - 85 yrs with: -- an active diagnosis of essential HTN between <u>01/01/2017 and 06/30/2017</u>	<ul style="list-style-type: none"> Active diagnosis through <i>start</i> of MP: <ul style="list-style-type: none"> -- ESRD -- Pregnancy -- Stage 5 CKD Diagnosed <i>before or during</i> MP: <ul style="list-style-type: none"> -- Dialysis -- History of renal transplant TENTATIVE - Patients ≥ 65 yrs in an Institutional SNP or residing in a long-term care facility Deceased during MP In hospice, including non-hospice with palliative goals or comfort care at anytime during MP Moved out of country at anytime during MP Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> CKD = Chronic Kidney Disease ESRD = End-Stage Renal Disease HMO = Health Maintenance Organization MP = Measurement Period = 01/01/2017 - 12/31/2017 MR = Medical Record SNP = Special Needs Plan WI = Web Interface 	<ol style="list-style-type: none"> Only the <i>most recent</i> BP reading that is performed by a clinician with appropriate skills and experience, <i>in the provider office</i>, are acceptable If there are multiple BP readings on the same day, use the <i>lowest systolic and diastolic</i> measurements (from different readings, if appropriate) as the most recent BP reading Patient-reported BP values are <i>not acceptable</i> BP screening <u>cannot</u> be completed during a telehealth encounter
IVD-2 Ischemic Vascular Disease (IVD) - Use of Aspirin or Another Antiplatelet	Patients ≥ 18 yrs who were: -- diagnosed with AMI, CABG, or PCI between 01/01/2016 and 12/31/2016 OR -- active diagnosis of IVD <i>during</i> the MP and -- had documentation of ASA or other antiplatelet use during the MP	<ul style="list-style-type: none"> Documentation of Denominator Exclusion Anticoagulant Medication use overlapping the MP Diagnosis of PVD and/or PAD do not qualify as IVD Deceased during MP In hospice, including non-hospice with palliative goals or comfort care at anytime during MP Moved out of country at anytime during MP Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> Active Diagnosis = a diagnosis that is: <ul style="list-style-type: none"> -- on the patient's problem list -- a diagnosis code on the encounter, or -- documented in a progress note indicating that the patient is being treated or managed for the disease or condition during the MP AMI = Acute Myocardial Infarction ASA = Aspirin CABG = Coronary Artery Bypass Graft HMO = Health Maintenance Organization IVD = Ischemic Vascular Disease MP = Measurement Period = 01/01/2017 - 12/31/2017 MR = Medical Record PAD = Peripheral Arterial Disease PCI = Percutaneous Coronary Intervention PVD = Peripheral Vascular Disease WI = Web Interface 	<ol style="list-style-type: none"> Denominator Exclusion Anticoagulant Medications: <ul style="list-style-type: none"> -- Apixaban -- Argatroban -- Bivalirudin -- Dabigatran -- Dalteparin -- Desirudin -- Edoxaban -- Enoxaparin -- Fondaparinux -- Heparin -- Lepirudin -- Rivaroxaban -- Tinzaparin -- Warfarin Oral antiplatelet meds include: <ul style="list-style-type: none"> -- ASA -- clopidogrel or combination of aspirin and extended-release dipyridamole -- Prasugrel -- Ticagrelor -- Ticlopidine Antiplatelet use documentation <u>can</u> be completed during a telehealth encounter
MH-1 Depression Remission at Twelve Months	Patients ≥ 18 yrs with: -- major depression or dysthymia AND -- an initial PHQ-9 > 9 documented between 12/1/2015 and 11/30/2016 AND -- demonstrate remission as evidenced by PHQ-9 < 5 at 12 months (+ or - 30 days) after index visit	<ul style="list-style-type: none"> Active diagnosis of bipolar disorder Active diagnosis of personality disorder Permanent Nursing Home Resident Deceased during MP In hospice, including non-hospice with palliative goals or comfort care at anytime during MP Moved out of country at anytime during MP Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> Active Diagnosis = a diagnosis that is: <ul style="list-style-type: none"> -- on the patient's problem list -- a diagnosis code on the encounter, or -- documented in a progress note indicating that the patient is being treated or managed for the disease or condition during the MP HMO = Health Maintenance Organization MP = Measurement Period = 01/01/2017-12/31/2017 MR = Medical Record Permanent Nursing Home Resident = a patient who is residing in a skilled nursing facility on a long-term basis. <ul style="list-style-type: none"> -- It does not include patients who are receiving short-term rehabilitative services following a hospital stay or patients residing in assisted-living or group-home settings WI = Web Interface 	<ol style="list-style-type: none"> Patient must be aged 18 years of age or older at Index Date Index PHQ-9 > 9 must be documented between 12/1/2015 and 11/30/2016; <ul style="list-style-type: none"> -- <i>if more than one, use most recent</i> PHQ-9 can be administered via telephone, email, mail, e-visit, patient portal, iPad/tablet, or patient kiosk Remission PHQ-9 < 5 must be documented during 12 months post-index date, +/- 30 days

2017 CMS Quality Reporting Quick Reference Guide

Measure	Description	Exclusions	Definitions	Data Guidance
PREV-5 Breast Cancer Screening	Patients 50 - 74 yrs who had: -- a mammogram screen for breast cancer between 10/01/2015 and 12/31/2017	<ul style="list-style-type: none"> Bilateral mastectomy 2 unilateral mastectomies TENTATIVE - Patients ≥ 65 yrs in an Institutional SNP or residing in a long-term care facility Deceased during MP In hospice, including non-hospice with palliative goals or comfort care at anytime during MP Moved out of country at anytime during MP Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> HMO = Health Maintenance Organization MP = Measurement Period = 01/01/2017 - 12/31/2017 MR = Medical Record SNP = Special Needs Plan WI = Web Interface 	<ol style="list-style-type: none"> MR documentation must include date the breast cancer screening was performed AND the result or findings Result of "normal" or "abnormal" is acceptable If patient reported, must include date, type of test, AND result/finding Testing includes: Breast x-ray, diagnostic mammography, mammogram, screening mammography, and 3D mammography <i>Not acceptable</i> are MRI and Ultrasound Documentation of screening mammography <u>can</u> be completed during a telehealth encounter Orders for breast cancer screening alone are <u>not</u> acceptable
PREV-6 Colorectal Cancer Screening	Patients 50 - 75 yrs who had <u>one</u> of the following: -- FOBT between 01/01/2017 and 12/31/2017 -- Flexible sigmoidoscopy between 01/01/2013 and 12/31/2017 -- Colonoscopy between 01/01/2008 and 12/31/2017 -- CT colonography between 01/01/2013 and 12/31/2017 -- FIT-DNA between 01/01/2015 and 12/31/2017	<ul style="list-style-type: none"> Diagnosis or past history of total colectomy or colorectal cancer TENTATIVE - Patients ≥ 65 yrs in an Institutional SNP or residing in a long-term care facility Deceased during MP In hospice, including non-hospice with palliative goals or comfort care at anytime during MP Moved out of country at anytime during MP Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> CT = Computed tomography DRE = Digital Rectal Exam FIT-DNA = Fecal immunochemical DNA test FOBT = Fecal Occult Blood Test HMO = Health Maintenance Organization MP = Measurement Period = 01/01/2017 - 12/31/2017 MR = Medical Record SNP = Special Needs Plan WI = Web Interface 	<ol style="list-style-type: none"> FOBT includes ColoCARE, Coloscreen, EZ Detect, FIT, flushable reagent pads, flushable reagent stool blood test, guiac smear test, Hemoccult, Seracult, stool occult blood test <i>Do not count</i> DRE or FOBT tests performed in the office <i>Do not count</i> FOBTs performed on a sample collected via DRE Documentation must include date of screening and result or findings -- Documentation of "normal" or "abnormal" is acceptable -- Patient Reported Requirement: date (year), type of test, and result/finding -- Documentation of colorectal cancer screening <u>can</u> be completed during a telehealth encounter
PREV-7 Influenza Immunization	Patients ≥ 6 mos who: -- received an influenza immunization during an office visit between 10/01/2016 and 03/31/2017 OR -- reported previously receiving an influenza immunization between 08/01/2016 and 03/31/2017	<ul style="list-style-type: none"> Medical reason, i.e., allergy Patient reason, i.e., refused System reason, i.e., vaccine unavailable Deceased during MP In hospice, including non-hospice with palliative goals or comfort care at anytime during MP Moved out of country at anytime during MP Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> HMO = Health Maintenance Organization MP = Measurement Period = 01/01/2017 - 12/31/2017 MR = Medical Record WI = Web Interface 	<ol style="list-style-type: none"> If WI is pre-filled with "Yes", no further action is required Documentation must include date of administration and type of vaccine administered Documented reason for not receiving influenza immunization: -- medical reason - e.g., patient allergy -- patient reason - e.g., patient declined -- system reason - e.g., vaccine unavailable Documentation of previously receiving the influenza immunization <u>can</u> be completed during a telehealth encounter
PREV-8 Pneumococcal Vaccination Status for Older Adults	Patients ≥ 65 yrs who: -- have ever received a pneumococcal vaccine	<ul style="list-style-type: none"> Deceased during MP In hospice, including non-hospice with palliative goals or comfort care at anytime during MP Moved out of country at anytime during MP Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> HMO = Health Maintenance Organization MP = Measurement Period = 01/01/2017 - 12/31/2017 MR = Medical Record WI = Web Interface 	<ol style="list-style-type: none"> Either PCV13, PPSV23 or both vaccines are acceptable Patient Reported Requirement: date (year) and type of vaccine Documentation of pneumococcal vaccination status <u>can</u> be completed during a telehealth encounter

2017 CMS Quality Reporting Quick Reference Guide

Measure	Description	Exclusions	Definitions	Data Guidance
<p>PREV-9 Body Mass Index (BMI) Screening and Follow-Up Plan</p>	<p>Patients ≥ 18 yrs with:</p> <ul style="list-style-type: none"> -- a documented BMI during the current encounter or during 6 mos prior to the current encounter <li style="text-align: center;">AND -- for BMI <i>above or below normal range</i> ≥ 18.5 and < 25 kg/m², a follow-up plan is documented during the encounter or during the 6 mos prior to the current encounter 	<ul style="list-style-type: none"> • Pregnancy • Receiving palliative care • Refusal of height/weight measurement or follow-up • Patients ≥ 65 yrs for whom weight loss/gain would complicate other underlying conditions, such as: <ul style="list-style-type: none"> -- Illness or physical disability -- Mental illness, dementia, confusion -- Nutritional or vitamin/mineral deficiency • Urgent or emergent patients where treatment delay would jeopardize patient health • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>BMI</i> - Body Mass Index • <i>Follow-Up Plan</i> - proposed outline of treatment for abnormal BMI • <i>Encounter</i> - patient visit when BMI is recorded • <i>HMO</i> = Health Maintenance Organization • <i>MP</i> = Measurement Period = 01/01/2017 - 12/31/2017 • <i>MR</i> = Medical Record • <i>WI</i> = Web Interface 	<p>(1) Follow-Up Plan may include documentation of:</p> <ul style="list-style-type: none"> -- Education -- Referral to a registered dietician, nutritionist, occupational therapy, physical therapy, PCP, exercise physiologist, mental health professional, or surgeon, e.g., "Patient referred to nutrition counseling for BMI above or below normal parameters." -- Pharmacological interventions -- Dietary supplements -- Exercise counseling -- Nutrition counseling <p>(2) Self-reported weights cannot be used.</p> <p>(3) Height and weight may be obtained from separate encounters</p> <p>(4) If > 1 BMI on chart, <i>use most recent</i></p> <p>(5) BMI calculation and recommended follow-up plan <u>cannot</u> be completed during a telehealth encounter</p>
<p>PREV-10 Tobacco Use - Screening and Cessation Intervention</p>	<p>Patients ≥ 18 yrs who were:</p> <ul style="list-style-type: none"> -- screened for tobacco use one or more times between 01/01/2016 and 12/31/2017 <li style="text-align: center;">AND -- who received cessation counseling intervention if identified as a tobacco user 	<ul style="list-style-type: none"> • Documentation of medical reason for <u>not screening</u> for tobacco use, such as: <ul style="list-style-type: none"> -- limited life expectancy -- other medical reason • Only medical reason excluding patient from receiving cessation intervention is limited life expectancy • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>HMO</i> = Health Maintenance Organization • <i>MP</i> = Measurement Period = 01/01/2017 - 12/31/2017 • <i>MR</i> = Medical Record • <i>Tobacco Use</i> = Includes any type of tobacco, e.g., smoke and smokeless • <i>Tobacco Cessation Intervention</i> = Includes brief counseling (≤ 3 mins) and/or pharmacotherapy • <i>WI</i> = Web Interface 	<p>(1) Confirm patient screened for tobacco at least once between 01/01/2016 and 12/31/2017</p> <p>(2) If more than one Tobacco Use Screening is on the chart, use most recent</p> <p>(3) Tobacco Use Screening and the Tobacco Cessation Intervention can occur at different encounters however they both must have occurred between 01/01/2016 and 12/31/2017</p> <p>(4) <i>Both</i> the Tobacco Screening <i>and</i> the Tobacco Cessation Intervention <u>can</u> be completed during a telehealth encounter</p> <p>(5) Electronic Nicotine Delivery Systems (ENDS) <u>are not considered evidence of tobacco use and cannot be used as a tobacco cessation aid</u></p>
<p>PREV-12 Screening for Depression and Follow-Up Plan</p>	<p>Patients ≥ 12 yrs who were:</p> <ul style="list-style-type: none"> -- screened for depression on the date of encounter using an age appropriate standardized depression screening tool <li style="text-align: center;">AND -- if positive, a follow-up plan is documented on the date of the positive screen 	<ul style="list-style-type: none"> • Active diagnosis of depression • Active diagnosis of bipolar disorder • Patient refusal to participate • Urgent or emergent patients where treatment delay would jeopardize patient health • Patient's functional capacity or motivation to improve may impact the accuracy of results of standardized depression assessment tools • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>Active Diagnosis</i> = a diagnosis that is: <ul style="list-style-type: none"> -- on the patient's problem list -- a diagnosis code on the encounter, or -- documented in a progress note indicating that the patient is being treated or managed for the disease or condition during the MP • <i>HMO</i> = Health Maintenance Organization • <i>MP</i> = Measurement Period = 01/01/2017 - 12/31/2017 • <i>MR</i> = Medical Record • <i>Standardized Depression Screening Tool</i> = a normalized and validated depression screening tool developed for the patient population in which it is being utilized • <i>WI</i> = Web Interface 	<p>(1) Adolescent Screening Tools (12 - 17 yrs) - see POM ACO website for suggestions</p> <p>(2) Adult Screening Tools (≥ 18 yrs) - see POM ACO website for suggestions</p> <p>(3) Name of the standardized depression screening tool used must be documented in the medical record</p> <p>(4) The depression screening must be reviewed and addressed in the office of the provider filing the code, on the date of the encounter</p> <p>(5) The screening and encounter must occur on the same date</p> <p>(6) Follow-Up Plan <u>must include one or more of the following</u>:</p> <ul style="list-style-type: none"> -- Additional evaluation for depression -- Suicide Risk Assessment -- Referral to a practitioner who is qualified to diagnose and treat depression -- Pharmacological interventions -- Other interventions or follow-up for the diagnosis or treatment of depression <p>(7) Follow-up plan must be related to the positive depression screening, e.g., "Patient referred for psychiatric evaluation due to positive depression screening."</p> <p>(8) Screening for depression <u>can</u> be completed during a telehealth encounter however the results <u>must</u> be reviewed/verified <u>and</u> documented by the eligible professional in the medical record <i>on the date of the encounter</i> to meet the screening portion of this measure</p> <p>(9) Documentation of recommended follow-up plan for a positive depression screen <u>can</u> be completed during a telehealth encounter</p>

2017 CMS Quality Reporting Quick Reference Guide

Measure	Description	Exclusions	Definitions	Data Guidance
PREV-13 Statin Therapy for the Prevention and Treatment of Cardiovascular Disease	<p>Patients ≥ 21 yrs at beginning of MP who:</p> <ul style="list-style-type: none"> -- have a history of or an active diagnosis of ASCVD <p style="text-align: center;">OR</p> <p>Patients ≥ 21 yrs at beginning of MP who:</p> <ul style="list-style-type: none"> -- have ever had a fasting or direct LDL-C ≥ 190 mg/dL or -- were previously diagnosed or have an active diagnosis of familial / pure hypercholesterolemia <p style="text-align: center;">OR</p> <p>Patients 40 - 75 yrs at beginning of MP with:</p> <ul style="list-style-type: none"> -- a diagnosis of Type 1 or 2 DM and a fasting or direct LDL-C of 70 - 189 mg/dL -- recorded as the highest result between 1/1/2015 through 12/31/2017 	<ul style="list-style-type: none"> • Active diagnosis of pregnancy • Breastfeeding • Diagnosis of rhabdomyolysis • History of adverse effect, allergy, or intolerance to statin medication • Palliative care • Active liver disease or hepatic disease or insufficiency • ESRD • DM diagnosis with most recent fasting or direct LDL-C < 70 mg/dL and are not taking statin therapy • Deceased during MP • In hospice, including non-hospice with palliative goals or comfort care at anytime during MP • Moved out of country at anytime during MP • Enrolled in HMO at any time during MP 	<ul style="list-style-type: none"> • <i>Active Diagnosis</i> = a diagnosis that is: <ul style="list-style-type: none"> -- on the patient's problem list -- a diagnosis code on the encounter, or -- documented in a progress note indicating that the patient is being treated or managed for the disease or condition during the MP • ASCVD = Clinical Atherosclerotic Cardiovascular Disease includes: <ul style="list-style-type: none"> -- Acute coronary syndromes -- History of myocardial infarction -- Stable or unstable angina -- Coronary or other arterial revascularization -- Stroke or transient ischemic attack (TIA) -- Peripheral arterial disease of atherosclerotic origin • DM = Diabetes mellitus • ESRD = End-stage renal disease • HMO = Health Maintenance Organization • LDL-C = Low-Density Lipoprotein Cholesterol • MP = Measurement Period = 01/01/2017 - 12/31/2017 • MR = Medical Record • WI = Web Interface 	<ol style="list-style-type: none"> (1) Adherence to statin therapy is not part of this measure (2) If laboratory unable to calculate LDL-C value due to high triglycerides, select "No". (3) If the test result is labeled "unreliable" and a result is provided, select "No" (4) Diabetes history is defined as any history of diabetes, prior to or during the measurement period (5) Documentation of statin therapy prescribed or being taken during the measurement period <u>cannot</u> be completed during a telehealth encounter